



Autumn Crocus



Crocus



Snow Drop



Oregon Iris



Grape Hyacinth



Daffodil



Common Yarrow



Pearly Everlasting



Sea Pink



Western Columbine

COMMON NAME SCIENTIFIC NAME	OREGON NATIVE	HEIGHT	SPREAD	LIGHT	EVERGREEN/DECIDUOUS	SEASONAL INTEREST	COLOR/FEATURE	WATER	MAINTENANCE	WILDLIFE	COMMENTS
Autumn Crocus <i>Colchicum autumnale</i>	N	1/2'	1'	sun	D	fall bloom is a bright welcome sight	white, pink, pale purple	L	little necessary, divide clumps in spring	N	many other varieties of this handsome fall blooming bulb
Crocus <i>Crocus spp.</i>	N	3-6"	1/2 - 1'	sun	D	early spring flowers	yellow, white, violet, and combinations	L	little necessary, divide clumps in autumn	N	many fine varieties are available, some may be naturalized in a lawn area
Snow Drop <i>Galanthus nivalis</i>	N	6-9"	3"	sun to part shade	D	early spring flowers	white flowers	L	little maintnace	N	plant in grouping for greatest effect
Oregon Iris <i>Iris tenax</i>	Y	1'	1'	sun	E	spring flower display	light purple or white flowers	L	trim back damaged foliage	N	slugs damage foliage, will reseed readily, other fine native iris are available
Grape Hyacinth <i>Muscari armeniacum</i>	N	1/2'	1/2'	sun	D	spring display of grape-like clusters of flowers	purplish flowers in clusters	L	trim back damaged foliage	N	Valerie Finnis' is a superb light blue, this bulb multiplies
Daffodil <i>Narcissus spp.</i>	N	6"-1 1/2'	6"	sun	D	early spring flowers	various shades of orange and yellow flowers	L	little maintenance	N	tremendous variety of flower forms and bloom period

Perennials

Common Yarrow <i>Achillea millifolium</i>	N	1 1/2-2'	1 1/2-2'	sun	D	summer flowers, fine textured foliage	many flower colors	L	cut off old flower stalks and dead foliage before spring growth begins	N	spreads by rhizomes, many varieties and colors from white, to orange, to red
Pearly Everlasting <i>Anaphalis margaritacea</i>	Y	3'	2'	sun	D	white long-lasting flowers	silver-downy foliage	L	cut off old flower stalks and dead foliage before spring growth begins	N	flowers may be dried and used in floral arrangements, may become weedy
Sea Pink <i>Armeria maitina</i>	N	1'	1'	sun	D	dense mound of fine textured foliage	white to pink flowers on 10 inch stems	M	remove damaged leaves and dead flowers	N	when planting do not plant to deep, do not mound bark dust to close to this plant
Western Columbine <i>Aquilegia formosa</i>	Y	2'	2'	sun	D	showy flowers attract humming birds	orange and yellow flowers	L	cut off old flower stalks and dead foliage before spring growth begins	Y	nodding flowers are 1-2" across, plant will reseed



Wormwood



Aster



False Indigo



Snow-In-Summer



Tickseed



Cheddar Pink



Western Bleeding Heart



Purple Cone Flower



California Fuchsia



Cushion Spurge

COMMON NAME SCIENTIFIC NAME	OREGON NATIVE	HEIGHT	SPREAD	LIGHT	EVERGREEN/DECIDUOUS	SEASONAL INTEREST	COLOR/FEATURE	WATER	MAINTENANCE	WILDLIFE	COMMENTS
Wormwood <i>Artemisia suksdorfii</i>	Y	1-2'	1-2'	sun	E	finely cut leaves	gray-green leaves	L	remove damaged foliage	N	several species of native <i>Artemisia</i>
Aster <i>Aster subspicatus</i>	Y	3'	2'	sun	D	summer flowers	purple flowers	L	cut off old flower stalks and dead foliage before spring growth begins	N	several species of native Aster, may spread
False Indigo <i>Baptisa australis</i>	N	3'	3-4'	sun	D	spring flowers and seed heads	light purple flowers	M	cut off old flower stalks and dead foliage before spring growth begins	N	seed pods dry and rattle in the breeze
Snow-In-Summer <i>Cerastium tomentosum</i>	N	8"	3'	sun to part shade	D	dense tufty mats of silvery gray	white flowers	M	tip shear after bloom is finished to have a more dense plant	N	use as a ground-cover on slopes, may be used in combination with bulbs
Tickseed <i>Coreopsis grandiflora</i>	N	18"	3'	sun	D	flowers bloom all summer	brightly yellow flowers	L	cut to ground level prior to new spring growth	N	spreads by self-sowing, several outstanding varieties available - many which will not self-sow, and many with attractive leaf forms
Cheddar Pink <i>Dianthus gratianopolitanus</i>	N	1'	1'	sun	E	fragrant flowers and mat-forming foliage	pink flowers, gray-green leaves	L	remove damaged foliage	N	tip shearing in spring may help to make clumps more dense
Western Bleeding Heart <i>Dicentra formosa</i>	Y	1 1/2'	3'	sun to part shade	D	stalks of pendulous pink flowers	blue-green leaves	L	cut to ground level prior to new spring growth	N	may be used as a ground-cover, spreads freely by under-ground stems, attractive foliage all summer in shade
Purple Cone Flower <i>Echinacea purpurea</i>	N	2-3'	2'	sun	D	blooms over a long period in summer	light purple flowers	M	cut to ground level prior to new spring growth	Y	brown seed heads may be dried and used in floral arrangements, birds eat seeds
California Fuchsia <i>Epilobium canum</i>	N	2'	4'	sun	D	many flowers in late summer and autumn	red flowers	L	cut to ground level prior to new spring growth	Y	see *, hummingbirds feed from nectar in flowers, often sold as <i>Zauschneria</i>
Cushion Spurge <i>Euphorbia polychroma</i>	N	1 1/2'	2'	sun to part shade	E	fall color, neatly rounded plants	yellow to orange-red leaves in fall	M	cut back damaged foliage	N	deep green leaves are symmetrically arranged, plants grown in full sun have better fall color

*Plants noted as weedy or invasive in other states or countries



Blanket Flower



Gaura



Cranesbill



Bloody Cranesbill



Corsican Hellebore



Candytuft



Red Hot Poker



Gayfeather



Blue Flax



Lupine

COMMON NAME SCIENTIFIC NAME	OREGON NATIVE	HEIGHT	SPREAD	LIGHT	EVERGREEN/DECIDUOUS	SEASONAL INTEREST	COLOR/FEATURE	WATER	MAINTENANCE	WILDLIFE	COMMENTS
Blanket Flower <i>Gaillardia x grandiflora</i>	N	1'	1'	sun	D	long bloom period during summer	red and yellow flowers with maroon bands	L	cut to ground level prior to spring growth and after flowering to encourage repeat bloom	N	many varieties available, flowers throughout summer
Gaura <i>Gaura lindheimeri</i>	N	2-3'	2-3'	sun	D	long bloom period during summer	white to light pink flowers	M	cut to ground level prior to new spring growth	N	see *, graceful feathery flower stalks, will happily re-seed in the garden
Cranesbill <i>Geranium macrorrhizum</i>	N	8-10"	1 1/2'	sun	D	blooms late spring through early summer	white, pink, or magenta flowers	L	cut to ground level prior to new spring growth	N	see *, attractive autumn foliage color - yellow to orange and scarlet
Bloody Cranesbill <i>Geranium sanguineum</i>	N	1 1/2'	2'	sun to part shade	D	dense mound with abundant flowers	white to pink	L	cut to ground level prior to new spring growth	N	plants grown in sun have foliage which turns red in fall, will re-bloom in mid summer if cut to ground
Corsican Hellebore <i>Helleborus argutifolius</i>	N	2-3'	3-4'	part shade to shade	E	flowers in winter	pale green flowers	M	remove damaged leaves and dead flowers	N	large bold texture, 6-9 inch leaves are bluish green, many varieties available
Candytuft <i>Iberis sempervirens</i>	N	1'	1-2'	sun to part shade	E	abundant flowers at branch tips	white flowers, tiny dark green leaves	L	tip shear after bloom is finished to have a more dense plant	N	plant is covered with flowers in spring
Red Hot Poker <i>Kniphofia uvaria</i>	N	2-4'	2-3'	sun to part shade	E	slender bold colored flowers arise above the plant	orange-red flowers	M	cut to ground level prior to new spring growth	Y	many varieties with other color forms available, flowers attract hummingbirds
Gayfeather <i>Liatris spicata</i>	N	2-3'	1 1/2'	sun	D	grassy clumps of leaves topped by fuzzy flower spikes	light purple	M	cut to ground level prior to new spring growth	N	other varieties with pink or white flowers are available
Blue Flax <i>Linum perenne</i>	Y	2'	1' - 1 1/2'	sun	D	feathery stems, flowers close in shade or late in the day	tiny light blue flowers	L	cut to ground level prior to new spring growth	N	self-sows, grown for fiber (linen) and seed (linseed oil)
Lupine <i>Lupinus</i>	Y	2-3'	2' - 2 1/2'	sun	D	large stalks of flowers in spring	leaves may be featured	M	cut to ground level prior to new spring growth	Y	see *, pastel colors to dark purple flowers, removing spent flowers encourages repeat bloom, many native species, self-sows readily

*Plants noted as weedy or invasive in other states or countries



Rose Mallow



Opium Poppy



Beardlip Penstemon




Russian Sage



Jerusalem Sage



Moss Phlox



Sword Fern



Prairie Coneflower



Matilija Poppy



Sage

COMMON NAME SCIENTIFIC NAME	OREGON NATIVE	HEIGHT	SPREAD	LIGHT	EVERGREEN/DECIDUOUS	SEASONAL INTEREST	COLOR/FEATURE	WATER	MAINTENANCE	WILDLIFE	COMMENTS
Rose Mallow <i>Malva alcea</i>	N	3-4'	2'	sun	D	sub-shrub that flowers abundantly	light to dark pink flowers	M	cut to ground level prior to new spring growth	N	used for perennial borders or edging, not a long-lived plant
Opium Poppy <i>Papaver somniferum</i>	N	3'	1'	sun	D	large showy flowers	many colors of flowers	L	shake seeds from pods once they have dried to 'reseed'	Y	while this specific Poppy is an annual - reseeding makes it beautifully reappear in the garden, blue green leaves
Beardlip Penstemon <i>Penstemon barbatus</i>	N	3'	2'	sun to part shade	D	flowers continue over a long period	spikes of red flowers	L	cut to ground level prior to new spring growth, tip shear to encourage density	N	varieties with pink flowers are available, many excellent native <i>Penstemon</i>
Russian Sage <i>Perovskia atriplicifolia</i>	N	3'	3'	sun	D	gray-green leaves with feathery appearance	light purple flowers	L	tip prune when young to develop a dense plant, do not cut back plant as new growth emerges from old stems	N	extremely resistant to heat and drought once established
Jerusalem Sage <i>Phlomis fruticosa</i>	N	2-3'	3-4'	sun to part shade	D	wooly grey/green leaves 6-8 inches long	golden yellow ball shaped flower clusters	M	cut back after flowering	N	upright flower stems continue to develop with tiers of flowers
Moss Phlox <i>Phlox subulata</i>	N	6" - 1'	1-2'	sun	E	dense mat-forming growth, needle-like evergreen leaves	white, pink or blue flowers	L	cut off old flower stalks and dead foliage before spring growth begins	N	spring flowers make sheets of color, attractive on slopes and in rock gardens, many other fine selections
Sword Fern <i>Polystichum munitum</i>	Y	3'	3'	sun to shade	E	leathery shiny fronds	dark green foliage	M	remove damaged leaves	Y	tolerates much shade, bold pacific northwest native
Prairie Coneflower <i>Ratibida columnifera</i>	N	2 1/2'	1'	sun	D	branched plants with deeply cut leaves	yellow to brownish purple flowers	L	cut to ground level prior to new spring growth	Y	may be used as a meadow plant, looks good with ornamental grasses
Matilija Poppy <i>Romneya coulteri</i>	N	4'	4'	sun	D	tissue paper flowers are produced over a long period	white flowers with yellow stamens	L	cut off old flower stalks and dead foliage before spring growth begins	N	spectacular flowers are long-lasting in floral arrangements
Sage <i>Salvia</i> spp.	N	1-3'	1-3'	sun	D	late summer and fall blooming	deep puple to red-rose colored flowers	L	shorten and shape plants prior to spring growth	Y	hummingbirds feed from nectar in flowers, many varieties available



Pincushion Flower



Oregon Stonecrop



Sedum



Brilliant Stonecrop



Goldenrod



Kinnikinnick



Beach Strawberry



Carpet Broom



St. Johnswort

COMMON NAME SCIENTIFIC NAME	OREGON NATIVE	HEIGHT	SPREAD	LIGHT	EVERGREEN/DECIDUOUS	SEASONAL INTEREST	COLOR/FEATURE	WATER	MAINTENANCE	WILDLIFE	COMMENTS
Pincushion Flower <i>Scabiosa</i>	N	1'-2'	1'-2'	sun	D	long bloom period during summer	white, pink or light blue flowers	L	cut off old flower stalks and dead foliage before spring growth begins	Y	excellent cut flowers, remove spent flowers to encourage abundant re-bloom
Oregon Stonecrop <i>Sedum oregonum</i>	Y	less than 6"	1/2-1'	sun	E	mat-forming plant	leaves turn bronze-purple in fall	M	shear off spent flowers	N	must have a well-drained site
Sedum <i>Sedum spathulifolium</i>	Y	6"	1' +	sun	E	forms large patches	short stems with clusters of yellow flowers	L	little maintenance needed for this plant	N	grows well in rocky areas and slopes with well drained soil
Brilliant Stonecrop <i>Hylotelephium spectabile</i>	N	1-1 1/2'	1-2'	sun	E	gray-green leaves with flower stems reaching upward	light-purple fall color	L	cut to ground level prior to new spring growth	N	flower stalks and seed heads provide fall and winter interest, remove flower stalks prior to spring growth
Goldenrod <i>Solidago</i> spp.	Y	1-3'	1-2'	sun	D	late summer bloom	bright yellow flowers	L	little maintenance is needed, few pests attack this plant	N	good for cut flowers
Ground Covers											
Kinnikinnick <i>Aretostaphylos uva-ursi</i>	Y	6-12"	1-3'	sun or part shade	E	shiny leaves, winter fruit	deep green leaves, orange-red fruit	M	tip-prune plants when young to encourage density, remove damaged stems	Y	fruit provides winter food for birds, site must have year-round drainage, other varieties are available
Beach Strawberry <i>Fragaria chiloensis</i>	Y	6-12"	1-2'	sun	D	small scale groundcover	white flowers in spring	L	trim off during late winter to encourage spring growth	Y	glossy dark green leaves have tooth-edged leaflets, produce a few seedy fruits
Carpet Broom <i>Genista pilosa</i>	N	1-1 1/2'	4-7'	sun	D	green stems with tiny leaves, dense ground-cover	bright yellow flowers	L	remove damaged stems, little maintenance needed	N	this low-growing plant is covered with bright yellow flowers in early spring, fairly fast growing
St. Johnswort <i>Hypericum calycinum</i>	N	1-1 1/2'	1'-2'	sun	E	showy summer flowers, blue-green foliage	yellow flowers	L	trim back or mow to ground level every 2-3 years or after severe winter cold	N	may be used as a ground-cover that is tolerant of heat, dense ground-cover, some plant parts are toxic